

# Elliptic operators on rough domains

José María Martell



AMS Spring Eastern Sectional Meeting

Special Session on Regularity Theory of PDEs  
and Calculus of Variations on Domains with Rough Boundaries

April 13, 2019

Hartford

## Section 1

### Introduction

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$
- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$
- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$

$$\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$$

- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$

- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$
- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$
- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Lipschitz domains

Theorem (Dahlberg 1977)

$$\Omega \subset \mathbb{R}^{n+1} \text{ Lipschitz domain} \implies \begin{cases} \omega \in RH_2(\sigma) \\ \left( \int_{\Delta} k^2 d\sigma \right)^{\frac{1}{2}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$


- $\omega \in RH_2(\sigma) \iff$  Solvability of  $(D_2)$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^2(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_2)$
- Can we go beyond Lipschitz domains?

$$\omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \rightsquigarrow (D_q) \text{ solvable for some } q \text{ (large)}$$

# Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} \text{ “=} f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball

$$\Delta(x, r) = B(x, r) \cap \partial\Omega, x \in \partial\Omega$$


- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$

## Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = "f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball

$$\Delta(x, r) = B(x, r) \cap \partial\Omega, x \in \partial\Omega$$


- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$

## Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = "f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball

$$\Delta(x, r) = B(x, r) \cap \partial\Omega, x \in \partial\Omega$$


- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$

## Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = "f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball

$$\Delta(x, r) = B(x, r) \cap \partial\Omega, \quad x \in \partial\Omega$$


- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$

## Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = "f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball

$$\Delta(x, r) = B(x, r) \cap \partial\Omega, \quad x \in \partial\Omega$$


- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$

## Harmonic measure

- $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 2$ , open
- Harmonic measure  $\{\omega^X\}_{X \in \Omega}$  family of “probabilities” on  $\partial\Omega$

$$u(X) = \int_{\partial\Omega} f(x) d\omega^X(x) \quad \text{solves} \quad (D) \quad \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = "f \in C_c(\partial\Omega) \end{cases}$$



- Surface ball


$$\Delta(x, r) = B(x, r) \cap \partial\Omega, \quad x \in \partial\Omega$$

- $\sigma = \mathcal{H}^n|_{\partial\Omega}$

- $\partial\Omega$  ADR  $\rightsquigarrow \sigma(\Delta(x, r)) \approx r^n, \quad x \in \partial\Omega$


## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition




## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition




## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition




## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition




## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition



## [Jerison-Kenig 1982] NTA/CAD domains


- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition



- $\Omega$  is CAD  $\equiv \left\{ \begin{array}{l} \bullet \text{Interior} \text{ Corkscrew and Harnack Chain} \\ \bullet \text{Exterior} \text{ Corkscrew} \\ \bullet \partial\Omega \text{ ADR} \end{array} \right.$

## [Jerison-Kenig 1982] NTA/CAD domains

- Openness  $\rightsquigarrow$  Corkscrew condition
- Path-connectedness  $\rightsquigarrow$  Harnack chain condition



- $\Omega$  is CAD  $\equiv \left\{ \begin{array}{l} \bullet \text{Interior} \text{ Corkscrew and Harnack Chain} \\ \bullet \text{Exterior} \text{ Corkscrew} \\ \bullet \partial\Omega \text{ ADR} \end{array} \right.$
- $\Omega$  is 1-sided CAD  $\equiv \left\{ \begin{array}{l} \bullet \text{Interior} \text{ Corkscrew and Harnack Chain} \\ \bullet \partial\Omega \text{ ADR} \end{array} \right.$

# NTA/CAD domains

Theorem (David-Jerison 1990; Semmes 1989)

$$\bullet \quad \Omega \subset \mathbb{R}^{n+1} \text{ CAD} \quad \Rightarrow \quad \begin{cases} \omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \\ \left( \int_{\Delta} k^p d\sigma \right)^{\frac{1}{p}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$$

- $\bullet$   $\omega \in RH_p(\sigma) \iff$  Solvability of  $(D_{p'})$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^{p'}(\sigma) \end{cases}$
- $\bullet$   $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_{p'})$
- $\bullet$  Can we consider operators with variable coefficients?

# NTA/CAD domains

Theorem (David-Jerison 1990; Semmes 1989)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD  $\implies \begin{cases} \omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \\ \left( \int_{\Delta} k^p d\sigma \right)^{\frac{1}{p}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$
- $\omega \in RH_p(\sigma) \iff$  Solvability of  $(D_{p'}) \begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^{p'}(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_{p'})$
- Can we consider operators with variable coefficients?

# NTA/CAD domains

Theorem (David-Jerison 1990; Semmes 1989)

$$\bullet \quad \Omega \subset \mathbb{R}^{n+1} \text{ CAD} \quad \Rightarrow \quad \left\{ \begin{array}{l} \omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \\ \left( \int_{\Delta} k^p d\sigma \right)^{\frac{1}{p}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{array} \right.$$

$$\bullet \quad \omega \in RH_p(\sigma) \iff \text{Solvability of } (D_{p'}) \quad \left\{ \begin{array}{l} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^{p'}(\sigma) \end{array} \right.$$

$$\bullet \quad u(X) = \int_{\partial\Omega} f(y) d\omega^X(y) \text{ solves } (D_{p'})$$

- Can we consider operators with variable coefficients?

# NTA/CAD domains

Theorem (David-Jerison 1990; Semmes 1989)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD  $\implies \begin{cases} \omega \in A_\infty(\sigma) = \bigcup_{p>1} RH_p(\sigma) \\ \left( \int_{\Delta} k^p d\sigma \right)^{\frac{1}{p}} \lesssim \int_{\Delta} k d\sigma = \frac{w(\Delta)}{\sigma(\Delta)} \end{cases}$
- $\omega \in RH_p(\sigma) \iff$  Solvability of  $(D_{p'})$   $\begin{cases} \mathcal{L}u = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^{p'}(\sigma) \end{cases}$
- $u(X) = \int_{\partial\Omega} f(y) d\omega^X(y)$  solves  $(D_{p'})$
- Can we consider operators with variable coefficients?

## Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
  - $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (symmetric)  

$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
  - $\omega_L$  elliptic measure

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)  

$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

Theorem (Hofmann-Martell; Delellis-Kenig-Pipher '01; Milicevic-Pipher-Toro '13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$
- $|\nabla A|\delta \in L^\infty(\Omega)$
- $|\nabla A|^2 \delta$  is a Carleson measure
- $\omega_{L_0} \in A_\infty(\sigma)$
- $A$  is a Carleson perturbation of  $A_0$
- $\omega_L \in A_\infty(\sigma)$
- Can we go beyond CAD?

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)
 
$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

Theorem (Hofmann-Martell; Delellis-Kenig-Pipher '01; Milicevic-Pipher-Toro '13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$ 
  - $|\nabla A|\delta \in L^\infty(\Omega)$
  - $|\nabla A|^2\delta$  is a Carleson measure
- $\omega_{L_0} \in A_\infty(\sigma)$
- $A$  is a Carleson perturbation of  $A_0$
- Can we go beyond CAD?

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)
 
$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

**Theorem** (Kenig-Pipher 01; Fefferman-Kenig-Pipher 91, Milakis-Pipher-Toro 13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$
- $\left. \begin{array}{l} |\nabla A| \delta \in L^\infty(\Omega) \\ |\nabla A|^2 \delta \text{ is a Carleson measure} \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$
- $\left. \begin{array}{l} \omega_{L_0} \in A_\infty(\sigma) \\ A \text{ is a Carleson perturbation of } A_0 \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$

- Can we go beyond CAD?

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)
 
$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

Theorem (Kenig-Pipher 01; Fefferman-Kenig-Pipher 91, Milakis-Pipher-Toro 13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$
- $\left. \begin{array}{l} |\nabla A| \delta \in L^\infty(\Omega) \\ |\nabla A|^2 \delta \text{ is a Carleson measure} \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$
- $\left. \begin{array}{l} \omega_{L_0} \in A_\infty(\sigma) \\ A \text{ is a Carleson perturbation of } A_0 \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$

- Can we go beyond CAD?

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)
 
$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

Theorem (Kenig-Pipher 01; Fefferman-Kenig-Pipher 91, Milakis-Pipher-Toro 13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$
- $\left. \begin{array}{l} |\nabla A| \delta \in L^\infty(\Omega) \\ |\nabla A|^2 \delta \text{ is a Carleson measure} \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$
- $\left. \begin{array}{l} \omega_{L_0} \in A_\infty(\sigma) \\ A \text{ is a Carleson perturbation of } A_0 \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$

- Can we go beyond CAD?

# Elliptic operators

- $Lu(X) = -\operatorname{div}(A\nabla u)(X), X \in \Omega$
- $A(X) = (a_{i,j}(X))_{1 \leq i,j \leq n+1}$  **real** (**symmetric**)  

$$A(X)\xi \cdot \xi \geq \Lambda^{-1}|\xi|^2 \quad \text{and} \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|$$
- $\omega_L$  elliptic measure

Theorem (Kenig-Pipher 01; Fefferman-Kenig-Pipher 91, Milakis-Pipher-Toro 13)

- $\Omega \subset \mathbb{R}^{n+1}$  CAD
- $\delta(X) := \operatorname{dist}(X, \partial\Omega)$
- $\left. \begin{array}{l} |\nabla A| \delta \in L^\infty(\Omega) \\ |\nabla A|^2 \delta \text{ is a Carleson measure} \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$
- $\left. \begin{array}{l} \omega_{L_0} \in A_\infty(\sigma) \\ A \text{ is a Carleson perturbation of } A_0 \end{array} \right\} \implies \omega_L \in A_\infty(\sigma)$
- Can we go beyond CAD?

## Section 2

### Characterizations of CAD

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}_0$
- ①  $\Omega$  CAD       $\iff$       ②  $\omega_L \in A_\infty(\sigma)$

- ①  $\implies$  ② [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian
- ②  $\implies$  ① [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$
- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
  - [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}_0$
- ①  $\Omega$  CAD       $\iff$       ②  $\omega_L \in A_\infty(\sigma)$

- ①  $\implies$  ② [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian
- ②  $\implies$  ① [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$
- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
  - [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}_0$
- ①  $\Omega$  CAD       $\iff$       ②  $\omega_L \in A_\infty(\sigma)$

- ①  $\implies$  ② [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian
- ②  $\implies$  ① [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$
- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
  - [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\begin{matrix} 1 \\ 1 \end{matrix} \Omega \text{ CAD} \iff \begin{matrix} 2 \\ 2 \end{matrix} \omega_L \in A_\infty(\sigma)$

$\begin{matrix} 1 \\ 1 \end{matrix} \implies \begin{matrix} 2 \\ 2 \end{matrix}$  [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian

$\begin{matrix} 2 \\ 2 \end{matrix} \implies \begin{matrix} 1 \\ 1 \end{matrix}$  [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$

- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
- [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\begin{matrix} 1 \\ 1 \end{matrix} \Omega \text{ CAD} \iff \begin{matrix} 2 \\ 2 \end{matrix} \omega_L \in A_\infty(\sigma)$

$\begin{matrix} 1 \\ 1 \end{matrix} \implies \begin{matrix} 2 \\ 2 \end{matrix}$  [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian

$\begin{matrix} 2 \\ 2 \end{matrix} \implies \begin{matrix} 1 \\ 1 \end{matrix}$  [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$

- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
- [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\begin{matrix} 1 \\ 1 \end{matrix} \Omega \text{ CAD} \iff \begin{matrix} 2 \\ 2 \end{matrix} \omega_L \in A_\infty(\sigma)$

- $\begin{matrix} 1 \\ 1 \end{matrix} \implies \begin{matrix} 2 \\ 2 \end{matrix}$  [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian
- $\begin{matrix} 2 \\ 2 \end{matrix} \implies \begin{matrix} 1 \\ 1 \end{matrix}$  [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$
- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
- [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure

# Elliptic operators

- $L = -\operatorname{div}(A\nabla) \in \mathbb{L}_0$ :
- $|\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|^2 \delta$  is a Carleson measure:

$$\sup_{\substack{x \in \partial\Omega \\ 0 < r < \operatorname{diam}(\partial\Omega)}} \frac{1}{\sigma(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} |\nabla A(Y)|^2 \delta(Y) dY < \infty$$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\begin{matrix} 1 \\ 1 \end{matrix} \Omega \text{ CAD} \iff \begin{matrix} 2 \\ 2 \end{matrix} \omega_L \in A_\infty(\sigma)$

- $\begin{matrix} 1 \\ 1 \end{matrix} \implies \begin{matrix} 2 \\ 2 \end{matrix}$  [Kenig-Pipher]  $\rightsquigarrow \mathbb{L}_0$ ; [David-Jerison; Semmes]  $\rightsquigarrow$  Laplacian
- $\begin{matrix} 2 \\ 2 \end{matrix} \implies \begin{matrix} 1 \\ 1 \end{matrix}$  [Hofmann-M.-Mayboroda-Toro-Zhao]  $\rightsquigarrow \mathbb{L}_0$
- [Hofmann-M.-Uriarte; Azzam-Hofmann-M.-Nyström-Toro]  $\rightsquigarrow$  Laplacian
- [Hofmann-M.-Toro]  $\rightsquigarrow |\nabla A| \delta \in L^\infty(\Omega) + |\nabla A|$  is a Carleson measure

# Elliptic operators

- $u(X) = \int_{\partial\Omega} f(y) d\omega_L^X(y), f \in C_c(\partial\Omega) \rightsquigarrow Lu = 0$
- $(D_p)$   $\begin{cases} Lu = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^p(\sigma) \\ \|Nu\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)} \end{cases}$
- $(D_p)$  solvable  $\iff \omega_L \in RH_{p'}(\sigma)$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$   $(D_p)$  is solvable for large  $p$

# Elliptic operators

- $u(X) = \int_{\partial\Omega} f(y) d\omega_L^X(y), f \in C_c(\partial\Omega) \rightsquigarrow Lu = 0$
- $(D_p)$   $\begin{cases} Lu = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^p(\sigma) \\ \|Nu\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)} \end{cases}$
- $(D_p)$  solvable  $\iff \omega_L \in RH_{p'}(\sigma)$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$   $(D_p)$  is solvable for large  $p$

# Elliptic operators

- $u(X) = \int_{\partial\Omega} f(y) d\omega_L^X(y), f \in C_c(\partial\Omega) \rightsquigarrow Lu = 0$
- $(D_p)$   $\begin{cases} Lu = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^p(\sigma) \\ \|Nu\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)} \end{cases}$
- $(D_p)$  solvable  $\iff \omega_L \in RH_{p'}(\sigma)$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$   $(D_p)$  is solvable for large  $p$

# Elliptic operators

- $u(X) = \int_{\partial\Omega} f(y) d\omega_L^X(y), f \in C_c(\partial\Omega) \rightsquigarrow Lu = 0$
- $(D_p)$   $\begin{cases} Lu = 0 \text{ in } \Omega \\ u|_{\partial\Omega} = f \in L^p(\sigma) \\ \|Nu\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)} \end{cases}$
- $(D_p)$  solvable  $\iff \omega_L \in RH_{p'}(\sigma)$

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$   $(D_p)$  is solvable for large  $p$

## Section 3

### Perturbation

# Motivation

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$  •  $\omega_L \in A_\infty \iff$  •  $(D_p)$  solvability  $p \gg 1$

- Can we consider other operators? (e.g., non-smooth coefficients)
- [Fefferman-Kenig-Pipher 1991, Milakis-Pipher-Toro 2013]  
 $A_\infty(\sigma)$  is stable under Carleson perturbation on CAD
- Goal: Develop perturbation theory on 1-sided CAD

# Motivation

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$   $\omega_L \in A_\infty \iff (D_p)$  solvability  $p \gg 1$

- Can we consider other operators? (e.g., non-smooth coefficients)
- [Fefferman-Kenig-Pipher 1991, Milakis-Pipher-Toro 2013]  
 $A_\infty(\sigma)$  is stable under Carleson perturbation on CAD
- Goal: Develop perturbation theory on 1-sided CAD

# Motivation

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$  •  $\omega_L \in A_\infty \iff$  •  $(D_p)$  solvability  $p \gg 1$

- Can we consider other operators? (e.g., non-smooth coefficients)
- [Fefferman-Kenig-Pipher 1991, Milakis-Pipher-Toro 2013]  
 $A_\infty(\sigma)$  is stable under Carleson perturbation on CAD
- Goal: Develop perturbation theory on 1-sided CAD

# Motivation

## Theorem

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
- $L \in \mathbb{L}_0$
- $\Omega$  CAD  $\iff$  •  $\omega_L \in A_\infty \iff$  •  $(D_p)$  solvability  $p \gg 1$

- Can we consider other operators? (e.g., non-smooth coefficients)
- [Fefferman-Kenig-Pipher 1991, Milakis-Pipher-Toro 2013]  
 $A_\infty(\sigma)$  is stable under Carleson perturbation on CAD
- **Goal:** Develop perturbation theory on 1-sided CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\begin{array}{c} \circ \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q \\ \Updownarrow \qquad q>1 \end{array}$$

•  $(D_q)$  is solvability  $q > 1$

$$\begin{array}{c} \circ \omega_{L_0} \in RH_p \\ \Updownarrow \qquad p' \end{array}$$

•  $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

• Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz      • Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

• Dahlberg  $\rightsquigarrow$  Lipschitz

• Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$
$$\Updownarrow$$

$$\bullet \omega_{L_0} \in RH_p$$
$$\Updownarrow$$

- $(D_q)$  is solvability  $q \gg 1$

- $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

- Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

- Dahlberg  $\rightsquigarrow$  Lipschitz

- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$
$$\Updownarrow$$

- ( $D_q$ ) is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$
$$\Updownarrow$$

- ( $D_{p'}$ ) is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

- Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

- Dahlberg  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$
$$\Updownarrow$$

•  $(D_q)$  is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$
$$\Updownarrow$$

•  $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

• Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz      • Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

• Dahlberg  $\rightsquigarrow$  Lipschitz      • Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$

$\Updownarrow$

•  $(D_q)$  is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$

$\Updownarrow$

•  $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

• Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz      • Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

• Dahlberg  $\rightsquigarrow$  Lipschitz      • Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$

$\Updownarrow$

- $(D_q)$  is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$

$\Updownarrow$

- $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

- Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

- Dahlberg  $\rightsquigarrow$  Lipschitz

- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$

$\Updownarrow$

- $(D_q)$  is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$

$\Updownarrow$

- $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

- Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

- Dahlberg  $\rightsquigarrow$  Lipschitz

- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

# Perturbation

- Good operator:  $L_0 u = -\operatorname{div}(A_0 \nabla u)$

$$\bullet \omega_{L_0} \in A_\infty = \bigcup_{q>1} RH_q$$
$$\Updownarrow$$

- $(D_q)$  is solvability  $q \gg 1$

$$\bullet \omega_{L_0} \in RH_p$$
$$\Updownarrow$$

- $(D_{p'})$  is solvable  $p'$

- Perturbed operator:  $Lu = -\operatorname{div}(A \nabla u)$

- Question 1: When  $\omega_L \in A_\infty$ ?

- Fefferman-Kenig-Pipher  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

- Question 2: When is  $\omega_L \in RH_p$ ?

- Dahlberg  $\rightsquigarrow$  Lipschitz
- Milakis-Pipher-Toro  $\rightsquigarrow$  CAD

## Large constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.; Cavero-Hofmann-M.-Toro)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| < \infty$

$$\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$$

$$(D_p) \text{ solvable for } p > p_0 \implies (D_p) \text{ solvable for } p > q_0$$

## Large constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.; Cavero-Hofmann-M.-Toro)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| < \infty$

$$\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$$

$$(D_p) \text{ solvable for } p > p_0 \implies (D_p) \text{ solvable for } p > q_0$$

## Large constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.; Cavero-Hofmann-M.-Toro)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| < \infty$

$$\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$$

$$(D_p) \text{ solvable for } p > p_0 \implies (D_p) \text{ solvable for } p > q_0$$

## Large constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.; Cavero-Hofmann-M.-Toro)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| < \infty$

$$\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$$

$$(D_p) \text{ solvable for } p > p_0 \implies (D_p) \text{ solvable for } p > q_0$$

## Large constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.; Cavero-Hofmann-M.-Toro)

- $\Omega$  1-sided CAD     •  $L_0, L$  elliptic operators     •  $\|\rho(A_0, A)\| < \infty$

$$\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$$

$$(D_p) \text{ solvable for } p > p_0 \implies (D_p) \text{ solvable for } p > q_0$$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- ①  $\Omega$  CAD       $\iff$       ②  $\omega_L \in A_\infty(\sigma)$

$$\boxed{\textcircled{2}} \implies \boxed{\textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega$  CAD

$$\boxed{\textcircled{1}} \implies \boxed{\textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega$  CAD  $\implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- ①  $\Omega$  CAD       $\iff$       ②  $\omega_L \in A_\infty(\sigma)$

$$\boxed{\textcircled{2}} \implies \boxed{\textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega$  CAD

$$\boxed{\textcircled{1}} \implies \boxed{\textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega$  CAD  $\implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- (1)  $\Omega$  CAD       $\iff$       (2)  $\omega_L \in A_\infty(\sigma)$

$$\boxed{\textcircled{2}} \implies \boxed{\textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega$  CAD

$$\boxed{\textcircled{1}} \implies \boxed{\textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega$  CAD  $\implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\textcircled{2} \implies \textcircled{1}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\textcircled{1} \implies \textcircled{2}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\boxed{\textcircled{2} \implies \textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\boxed{\textcircled{1} \implies \textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\boxed{\textcircled{2} \implies \textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\boxed{\textcircled{1} \implies \textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\boxed{\textcircled{2} \implies \textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\boxed{\textcircled{1} \implies \textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\boxed{\textcircled{2} \implies \textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\boxed{\textcircled{1} \implies \textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Large constant perturbation and Characterization of CAD

- $L_0 = -\operatorname{div}(A_0 \nabla) \in \mathbb{L}_0$ :  $|\nabla A_0| \delta \in L^\infty(\Omega) + |\nabla A_0|^2 \delta$  Carleson meas.
- $L = -\operatorname{div}(A \nabla) \in \mathbb{L}$ :  $\|\rho(A_0, A)\| < \infty$  for some  $L_0 \in \mathbb{L}_0$

## Corollary

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided CAD
  - $L \in \mathbb{L}$
- $$\textcircled{1} \quad \Omega \text{ CAD} \quad \iff \quad \textcircled{2} \quad \omega_L \in A_\infty(\sigma)$$

$$\boxed{\textcircled{2} \implies \textcircled{1}}$$

- [Cavero-Hofmann-M.]:  $\omega_L \in A_\infty(\sigma) \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Hofmann-M.-Mayboroda-Toro-Zhao]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \Omega \text{ CAD}$

$$\boxed{\textcircled{1} \implies \textcircled{2}}$$

- [Kenig-Pipher]:  $\Omega \text{ CAD} \implies \omega_{L_0} \in A_\infty(\sigma)$
- [Milakis-Pipher-Toro]:  $\omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma)$

## Small constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| \ll 1$

$$\omega_{L_0} \in RH_q(\sigma) \implies \omega_L \in RH_q(\sigma)$$

$$(D_p) \text{ solvable for } p \geq q' \implies (D_p) \text{ solvable for } p \geq q'$$

## Small constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| \ll 1$

$$\omega_{L_0} \in RH_q(\sigma) \implies \omega_L \in RH_q(\sigma)$$

$$(D_p) \text{ solvable for } p \geq q' \implies (D_p) \text{ solvable for } p \geq q'$$

## Small constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.)

- $\Omega$  1-sided CAD
- $L_0, L$  elliptic operators
- $\|\rho(A_0, A)\| \ll 1$

$$\omega_{L_0} \in RH_q(\sigma) \implies \omega_L \in RH_q(\sigma)$$

$$(D_p) \text{ solvable for } p \geq q' \implies (D_p) \text{ solvable for } p \geq q'$$

## Small constant perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\| = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\sigma(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 \delta(Y) dY$$

Theorem (Cavero-Hofmann-M.)

- $\Omega$  1-sided CAD     •  $L_0, L$  elliptic operators     •  $\|\rho(A_0, A)\| \ll 1$

$$\omega_{L_0} \in RH_q(\sigma) \implies \omega_L \in RH_q(\sigma)$$

$$(D_p) \text{ solvable for } p \geq q' \implies (D_p) \text{ solvable for } p \geq q'$$

## Section 4

### Other $A_\infty$ properties

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
  - Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
  - [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY < 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$
This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$
- $$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
  - Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
  - [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY < 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
  - Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
  - [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
  - Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
  - [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
- Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
- [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
  - Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
  - [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

# Motivation

- 1-sided CAD  $\rightsquigarrow \sigma = H^n|_{\partial\Omega}$  good measure  $\rightsquigarrow \omega_L \in A_\infty(\sigma)$ 
  - What happens if  $H^n|_{\partial\Omega}$  is a bad object?
- $\Omega$  a 1-sided CAD and  $\sup_{\Delta \subset \partial\Omega} \frac{1}{\sigma(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 \delta dY < \infty$ 
$$\left( \omega_{L_0} \in A_\infty(\sigma) \implies \omega_L \in A_\infty(\sigma) \right) \implies \omega_L \in A_\infty(\omega_{L_0})$$
- Can we directly prove  $\omega_L \in A_\infty(\omega_{L_0})$ ?
- [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD
$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_L \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

This condition does not involve  $\sigma$ !!!

## Main questions

- [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD

$$\sup_{\Delta \subset \partial \Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_{L_0} \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

- Question 1: Do we really need  $\partial\Omega$  ADR?
  - Good PDE background  $\rightsquigarrow \omega_{L_0}$  doubling, CFMS ...
- Question 2: Large constant case?

## Main questions

- [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD

$$\sup_{\Delta \subset \partial \Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_{L_0} \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

- Question 1: Do we really need  $\partial\Omega$  ADR?
  - Good PDE background  $\rightsquigarrow \omega_{L_0}$  doubling, CFMS ...
- Question 2: Large constant case?

## Main questions

- [Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]  $\rightsquigarrow \Omega$  CAD

$$\sup_{\Delta \subset \partial\Omega} \frac{1}{\omega_{L_0}(\Delta)} \iint_{B \cap \Omega} \rho(A_0, A)^2 G_{L_0} dY \ll 1 \implies \omega_{L_0} \in RH_2(\omega_{L_0}) \subset A_\infty(\omega_{L_0})$$

- **Question 1:** Do we really need  $\partial\Omega$  ADR?
  - Good PDE background  $\rightsquigarrow \omega_{L_0}$  doubling, CFMS ...
- **Question 2:** Large constant case?

## 1-sided NTA and CDC

- $\Omega \subset \mathbb{R}^{n+1}$  **1-sided NTA** (aka uniform domain)
  - Interior Corkscrew
  - Interior Harnack Chain
- $\Omega$  satisfies **CDC** (capacity density condition, aka uniform 2-fat)

$$\frac{\text{Cap}_2(\overline{B} \setminus \Omega, 2B)}{\text{Cap}_2(\overline{B}, 2B)} \gtrsim 1$$

where  $\text{Cap}_2(K, D) = \inf \left\{ \iint_D |\nabla v|^2 dX : v \in C_c^\infty(D), v \geq 1 \text{ in } K \right\}$

- **CDC** is a quantitative version of **Wiener regularity**
- **Examples:**
  - $\Omega$  NTA  $\implies$  CDC  $\rightsquigarrow \overline{B}' \subset \overline{B} \setminus \overline{\Omega} \subset \overline{B} \setminus \Omega$  (exterior)
  - 1-sided  $\Omega$  CAD  $\implies$  CDC  $\rightsquigarrow \overline{\Delta} \subset \overline{B} \cap \partial\Omega \subset \overline{B} \setminus \Omega$  (boundary)

## 1-sided NTA and CDC

- $\Omega \subset \mathbb{R}^{n+1}$  **1-sided NTA** (aka uniform domain)
  - Interior Corkscrew
  - Interior Harnack Chain
- $\Omega$  satisfies **CDC** (capacity density condition, aka uniform 2-fat)

$$\frac{\text{Cap}_2(\overline{B} \setminus \Omega, 2B)}{\text{Cap}_2(\overline{B}, 2B)} \gtrsim 1$$

where  $\text{Cap}_2(K, D) = \inf \left\{ \iint_D |\nabla v|^2 dX : v \in C_c^\infty(D), v \geq 1 \text{ in } K \right\}$

- **CDC** is a quantitative version of **Wiener regularity**
- **Examples:**
  - $\Omega$  NTA  $\implies$  CDC  $\rightsquigarrow \overline{B}' \subset \overline{B} \setminus \overline{\Omega} \subset \overline{B} \setminus \Omega$  (exterior)
  - 1-sided  $\Omega$  CAD  $\implies$  CDC  $\rightsquigarrow \overline{\Delta} \subset \overline{B} \cap \partial\Omega \subset \overline{B} \setminus \Omega$  (boundary)

## 1-sided NTA and CDC

- $\Omega \subset \mathbb{R}^{n+1}$  **1-sided NTA** (aka uniform domain)
  - Interior Corkscrew
  - Interior Harnack Chain
- $\Omega$  satisfies **CDC** (capacity density condition, aka uniform 2-fat)

$$\frac{\text{Cap}_2(\overline{B} \setminus \Omega, 2B)}{\text{Cap}_2(\overline{B}, 2B)} \gtrsim 1$$

where  $\text{Cap}_2(K, D) = \inf \left\{ \iint_D |\nabla v|^2 dX : v \in C_c^\infty(D), v \geq 1 \text{ in } K \right\}$

- **CDC** is a quantitative version of **Wiener regularity**
- Examples:
  - $\Omega$  NTA  $\implies$  CDC  $\rightsquigarrow \overline{B}' \subset \overline{B} \setminus \overline{\Omega} \subset \overline{B} \setminus \Omega$  (exterior)
  - 1-sided  $\Omega$  CAD  $\implies$  CDC  $\rightsquigarrow \overline{\Delta} \subset \overline{B} \cap \partial\Omega \subset \overline{B} \setminus \Omega$  (boundary)

## 1-sided NTA and CDC

- $\Omega \subset \mathbb{R}^{n+1}$  **1-sided NTA** (aka uniform domain)
  - Interior Corkscrew
  - Interior Harnack Chain
- $\Omega$  satisfies **CDC** (capacity density condition, aka uniform 2-fat)

$$\frac{\text{Cap}_2(\overline{B} \setminus \Omega, 2B)}{\text{Cap}_2(\overline{B}, 2B)} \gtrsim 1$$

where  $\text{Cap}_2(K, D) = \inf \left\{ \iint_D |\nabla v|^2 dX : v \in C_c^\infty(D), v \geq 1 \text{ in } K \right\}$

- **CDC** is a quantitative version of **Wiener regularity**
- **Examples:**
  - $\Omega$  NTA  $\implies$  CDC  $\rightsquigarrow \overline{B'} \subset \overline{B} \setminus \overline{\Omega} \subset \overline{B} \setminus \Omega$  (exterior)
  - 1-sided  $\Omega$  CAD  $\implies$  CDC  $\rightsquigarrow \overline{\Delta} \subset \overline{B} \cap \partial\Omega \subset \overline{B} \setminus \Omega$  (boundary)

## 1-sided NTA and CDC

- $\Omega \subset \mathbb{R}^{n+1}$  1-sided NTA (aka uniform domain)
  - Interior Corkscrew
  - Interior Harnack Chain
- $\Omega$  satisfies CDC (capacity density condition, aka uniform 2-fat)

$$\frac{\text{Cap}_2(\overline{B} \setminus \Omega, 2B)}{\text{Cap}_2(\overline{B}, 2B)} \gtrsim 1$$

where  $\text{Cap}_2(K, D) = \inf \left\{ \iint_D |\nabla v|^2 dX : v \in C_c^\infty(D), v \geq 1 \text{ in } K \right\}$

- CDC is a quantitative version of Wiener regularity
- Examples:
  - $\Omega$  NTA  $\implies$  CDC  $\rightsquigarrow \overline{B'} \subset \overline{B} \setminus \overline{\Omega} \subset \overline{B} \setminus \Omega$  (exterior)
  - 1-sided  $\Omega$  CAD  $\implies$  CDC  $\rightsquigarrow \overline{\Delta} \subset \overline{B} \cap \partial\Omega \subset \overline{B} \setminus \Omega$  (boundary)

# Perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} = \sup_{\Delta(x,r) \subset \partial\Omega} \frac{1}{\omega_0(\Delta(x,r))} \iint_{B(x,r) \cap \Omega} \rho(A_0, A)(Y)^2 G_{L_0}(Y) dY$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\rho(A_0, A)\|_{\omega_{L_0}} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} = \sup_{\Delta(x, r) \subset \partial\Omega} \frac{1}{\omega_0(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} \rho(A_0, A)(Y)^2 G_{L_0}(Y) dY$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\rho(A_0, A)\|_{\omega_{L_0}} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} = \sup_{\Delta(x, r) \subset \partial\Omega} \frac{1}{\omega_0(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} \rho(A_0, A)(Y)^2 G_{L_0}(Y) dY$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\rho(A_0, A)\|_{\omega_{L_0}} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} = \sup_{\Delta(x, r) \subset \partial\Omega} \frac{1}{\omega_0(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} \rho(A_0, A)(Y)^2 G_{L_0}(Y) dY$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\rho(A_0, A)\|_{\omega_{L_0}} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Disagreement between  $L_0$  and  $L$

$$\rho(A_0, A)(Y) := \sup_{Z \in B(Y, \frac{\delta(Y)}{2})} \frac{|A_0(Z) - A(Z)|}{\delta(Z)}$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} = \sup_{\Delta(x, r) \subset \partial\Omega} \frac{1}{\omega_0(\Delta(x, r))} \iint_{B(x, r) \cap \Omega} \rho(A_0, A)(Y)^2 G_{L_0}(Y) dY$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\rho(A_0, A)\|_{\omega_{L_0}} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\rho(A_0, A)\|_{\omega_{L_0}} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Conical square function

$$\mathcal{A}(\rho(A_0, A))(x) := \left( \iint_{\Gamma(x)} \frac{\rho(A_0, A)(Y)^2}{\delta(Y)^{n-1}} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Small constant case in the unit ball, Lipschitz, CAD  
[Fefferman; Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]
- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Conical square function

$$\mathcal{A}(\rho(A_0, A))(x) := \left( \iint_{\Gamma(x)} \frac{\rho(A_0, A)(Y)^2}{\delta(Y)^{n-1}} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Small constant case in the unit ball, Lipschitz, CAD  
[Fefferman; Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]
- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Conical square function

$$\mathcal{A}(\rho(A_0, A))(x) := \left( \iint_{\Gamma(x)} \frac{\rho(A_0, A)(Y)^2}{\delta(Y)^{n-1}} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Small constant case in the unit ball, Lipschitz, CAD  
[Fefferman; Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]
- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Conical square function

$$\mathcal{A}(\rho(A_0, A))(x) := \left( \iint_{\Gamma(x)} \frac{\rho(A_0, A)(Y)^2}{\delta(Y)^{n-1}} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Small constant case in the unit ball, Lipschitz, CAD  
[Fefferman; Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]
- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

# Perturbation

- Conical square function

$$\mathcal{A}(\rho(A_0, A))(x) := \left( \iint_{\Gamma(x)} \frac{\rho(A_0, A)(Y)^2}{\delta(Y)^{n-1}} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega$$

Theorem (Akman, Hofmann, M., Toro)

- $\Omega$  1-sided NTA
- $\Omega$  CDC
- $L_0, L$  elliptic operators

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} < \infty \implies \omega_L \in A_\infty(\omega_{L_0})$$

$$\|\mathcal{A}(\rho(A_0, A))\|_{L^\infty(\omega_{L_0})} \ll_p 1 \implies \omega_L \in RH_p(\omega_{L_0})$$

- Small constant case in the unit ball, Lipschitz, CAD  
[Fefferman; Fefferman-Kenig-Pipher; Milakis-Pipher-Toro]
- Large constant case is new even in  $\mathbb{R}_+^{n+1}$ , unit ball, Lipschitz ...

Thank you for your attention!!!